

Absolute Constants and Hyperbolic Trigonometry



www.pgadey.ca
@pgadey

- ▶ Isomorphism of the Cayley-Klein and Poincaré disk models
- ▶ Isomorphism of the two Poincaré models
- ▶ Hyperbolic trigonometry functions
- ▶ Absolute constants





UNIVERSITY OF
TORONTO
MISSISSAUGA

Cayley and Poincaré



www.pgadey.ca
@pgadey




UNIVERSITY OF
TORONTO
MISSISSAUGA

Cayley and Poincaré



www.pgadey.ca
@pgadey



<https://www.geogebra.org/3d/gzzatbga>




UNIVERSITY OF
TORONTO
MISSISSAUGA

Cayley and Poincaré



www.pgadey.ca
@pgadey



The chord XY is a geodesic in the Cayley-Klein model.




UNIVERSITY OF
TORONTO
MISSISSAUGA

<https://www.geogebra.org/3d/gzzatbga>

Cayley and Poincaré



www.pgadey.ca
@pgadey



The chord XY is a geodesic in the Cayley-Klein model.

The arc $XA'Y$ is the corresponding geodesic in the Poincaré model.

<https://www.geogebra.org/3d/gzzatbga>



UNIVERSITY OF
TORONTO
MISSISSAUGA

The Poincaré Disk and Half-Plane

Consider the map $\Omega : \mathbb{D}^2 \rightarrow \mathbb{C}_+$.

$$\Omega(z) = i \cdot \frac{1+z}{1-z}$$

www.pgadey.ca
@pgadey





UNIVERSITY OF
TORONTO
MISSISSAUGA

The Poincaré Disk and Half-Plane

Consider the map $\Omega : \mathbb{D}^2 \rightarrow \mathbb{C}_+$.

$$\Omega(z) = i \cdot \frac{1+z}{1-z}$$

www.pgadey.ca
@pgadey





UNIVERSITY OF
TORONTO
MISSISSAUGA

The Poincaré Disk and Half-Plane

Consider the map $\Omega : \mathbb{D}^2 \rightarrow \mathbb{C}_+$.

$$\Omega(z) = i \cdot \frac{1+z}{1-z}$$

www.pgadey.ca
@pgadey



It induces an isomorphism of geometries:





UNIVERSITY OF
TORONTO
MISSISSAUGA

The Poincaré Disk and Half-Plane

Consider the map $\Omega : \mathbb{D}^2 \rightarrow \mathbb{C}_+$.

$$\Omega(z) = i \cdot \frac{1+z}{1-z}$$

www.pgadey.ca
@pgadey



It induces an isomorphism of geometries:

$$(\mathbb{H}^2, \mathcal{M}) \simeq (\mathbb{C}_+, \mathbb{R}\text{Möb})$$



UNIVERSITY OF
TORONTO
MISSISSAUGA

The Hyperbolic Model Isomorphism Theorem

Corollary (10.2.3)

The three models of hyperbolic geometry, namely the Poincaré disk and half-plane models and the Cayley-Klein model, are isomorphic as geometries.

www.pgadey.ca
@pgadey



UNIVERSITY OF
TORONTO
MISSISSAUGA

Hyperbolic Trigonometry



www.pgadey.ca
@pgadey



UNIVERSITY OF
TORONTO
MISSISSAUGA

Hyperbolic Trigonometry

The hyperbolic trigonometric functions are defined by:

www.pgadey.ca
@pgadey



UNIVERSITY OF
TORONTO
MISSISSAUGA

Hyperbolic Trigonometry

The hyperbolic trigonometric functions are defined by:

$$\sinh(t) = \frac{e^t - e^{-t}}{2}$$

www.pgadey.ca
@pgadey



UNIVERSITY OF
TORONTO
MISSISSAUGA

Hyperbolic Trigonometry

The hyperbolic trigonometric functions are defined by:

$$\sinh(t) = \frac{e^t - e^{-t}}{2} \quad \cosh(t) = \frac{e^t + e^{-t}}{2}$$

www.pgadey.ca
@pgadey



UNIVERSITY OF
TORONTO
MISSISSAUGA

Hyperbolic Trigonometry

The hyperbolic trigonometric functions are defined by:

$$\sinh(t) = \frac{e^t - e^{-t}}{2} \quad \cosh(t) = \frac{e^t + e^{-t}}{2}$$

www.pgadey.ca
@pgadey

Theorem



UNIVERSITY OF
TORONTO
MISSISSAUGA

Hyperbolic Trigonometry

The hyperbolic trigonometric functions are defined by:

$$\sinh(t) = \frac{e^t - e^{-t}}{2} \quad \cosh(t) = \frac{e^t + e^{-t}}{2}$$

www.pgadey.ca
@pgadey

Theorem

For a hyperbolic triangle ABC with sides $\{a, b, c\}$ and angles $\{\alpha, \beta, \gamma\}$ we have:



UNIVERSITY OF
TORONTO
MISSISSAUGA

Hyperbolic Trigonometry

The hyperbolic trigonometric functions are defined by:

$$\sinh(t) = \frac{e^t - e^{-t}}{2} \quad \cosh(t) = \frac{e^t + e^{-t}}{2}$$

www.pgadey.ca
@pgadey

Theorem

For a hyperbolic triangle ABC with sides $\{a, b, c\}$ and angles $\{\alpha, \beta, \gamma\}$ we have:

$$\frac{\sinh(a)}{\sin(\alpha)}$$



UNIVERSITY OF
TORONTO
MISSISSAUGA

Hyperbolic Trigonometry

The hyperbolic trigonometric functions are defined by:

$$\sinh(t) = \frac{e^t - e^{-t}}{2} \quad \cosh(t) = \frac{e^t + e^{-t}}{2}$$

www.pgadey.ca
@pgadey

Theorem

For a hyperbolic triangle ABC with sides $\{a, b, c\}$ and angles $\{\alpha, \beta, \gamma\}$ we have:

$$\frac{\sinh(a)}{\sin(\alpha)} = \frac{\sinh(b)}{\sin(\beta)}$$



UNIVERSITY OF
TORONTO
MISSISSAUGA

Hyperbolic Trigonometry

The hyperbolic trigonometric functions are defined by:

$$\sinh(t) = \frac{e^t - e^{-t}}{2} \quad \cosh(t) = \frac{e^t + e^{-t}}{2}$$

www.pgadey.ca
@pgadey

Theorem

For a hyperbolic triangle ABC with sides $\{a, b, c\}$ and angles $\{\alpha, \beta, \gamma\}$ we have:

$$\frac{\sinh(a)}{\sin(\alpha)} = \frac{\sinh(b)}{\sin(\beta)} = \frac{\sinh(c)}{\sin(\gamma)}$$



UNIVERSITY OF
TORONTO
MISSISSAUGA

Absolute Constants



www.pgadey.ca
@pgadey



UNIVERSITY OF
TORONTO
MISSISSAUGA

Absolute Constants

In Euclidean space, there is no “natural” unit of measurement.

www.pgadey.ca
@pgadey



UNIVERSITY OF
TORONTO
MISSISSAUGA

Absolute Constants

In Euclidean space, there is no “natural” unit of measurement.
Similarity of figures implies there is no natural reference point.


www.pgadey.ca
@pgadey



UNIVERSITY OF
TORONTO
MISSISSAUGA

Absolute Constants

In Euclidean space, there is no “natural” unit of measurement.
Similarity of figures implies there is no natural reference point.




www.pgadey.ca
@pgadey



UNIVERSITY OF
TORONTO
MISSISSAUGA

Absolute Constants

In Euclidean space, there is no “natural” unit of measurement.
Similarity of figures implies there is no natural reference point.



www.pgadey.ca
@pgadey



UNIVERSITY OF
TORONTO
MISSISSAUGA

In hyperbolic geometry, there are natural units!

The Ideal Triangle



www.pgadey.ca
@pgadey



UNIVERSITY OF
TORONTO
MISSISSAUGA

The Ideal Triangle

We know that the area of a hyperbolic triangle is:

www.pgadey.ca
@pgadey



UNIVERSITY OF
TORONTO
MISSISSAUGA

The Ideal Triangle

We know that the area of a hyperbolic triangle is:

$$S = \pi - \alpha - \beta - \gamma$$

www.pgadey.ca
@pgadey



UNIVERSITY OF
TORONTO
MISSISSAUGA

The Ideal Triangle

We know that the area of a hyperbolic triangle is:

$$S = \pi - \alpha - \beta - \gamma \Rightarrow S \leq \pi$$

www.pgadey.ca
@pgadey



UNIVERSITY OF
TORONTO
MISSISSAUGA

The Ideal Triangle

We know that the area of a hyperbolic triangle is:

$$S = \pi - \alpha - \beta - \gamma \Rightarrow S \leq \pi$$

Thus, the largest triangle in hyperbolic space has area $S = \pi$.

www.pgadey.ca
@pgadey




UNIVERSITY OF
TORONTO
MISSISSAUGA

The Ideal Triangle

We know that the area of a hyperbolic triangle is:

$$S = \pi - \alpha - \beta - \gamma \Rightarrow S \leq \pi$$

Thus, the largest triangle in hyperbolic space has area $S = \pi$.



www.pgadey.ca
@pgadey




UNIVERSITY OF
TORONTO
MISSISSAUGA

The Ideal Triangle

We know that the area of a hyperbolic triangle is:

$$S = \pi - \alpha - \beta - \gamma \Rightarrow S \leq \pi$$

Thus, the largest triangle in hyperbolic space has area $S = \pi$.



The natural unit of measure for area is π .

www.pgadey.ca
@pgadey



UNIVERSITY OF
TORONTO
MISSISSAUGA

The Schweikart Radius



www.pgadey.ca
@pgadey



UNIVERSITY OF
TORONTO
MISSISSAUGA

The Schweikart Radius

Definition

www.pgadey.ca
@pgadey



UNIVERSITY OF
TORONTO
MISSISSAUGA

The Schweikart Radius

Definition

Given a pair of perpendicular rays OA and OB they meet the absolute \mathbb{A} at points X and Y .

www.pgadey.ca
@pgadey



UNIVERSITY OF
TORONTO
MISSISSAUGA

The Schweikart Radius

Definition

Given a pair of perpendicular rays OA and OB they meet the absolute \mathbb{A} at points X and Y . There is a unique line XY passing through X and Y .

www.pgadey.ca
@pgadey



UNIVERSITY OF
TORONTO
MISSISSAUGA

The Schweikart Radius

Definition

Given a pair of perpendicular rays OA and OB they meet the absolute \mathbb{A} at points X and Y . There is a unique line XY passing through X and Y . The distance from O to XY is the *Schweikart radius* denoted σ .

www.pgadey.ca
@pgadey




UNIVERSITY OF
TORONTO
MISSISSAUGA

The Schweikart Radius

Definition

Given a pair of perpendicular rays OA and OB they meet the absolute \mathbb{A} at points X and Y . There is a unique line XY passing through X and Y . The distance from O to XY is the *Schweikart radius* denoted σ .



www.pgadey.ca
@pgadey




UNIVERSITY OF
TORONTO
MISSISSAUGA

The Schweikart Radius

Definition

Given a pair of perpendicular rays OA and OB they meet the absolute \mathbb{A} at points X and Y . There is a unique line XY passing through X and Y . The distance from O to XY is the *Schweikart radius* denoted σ .



The natural unit of measure for length is σ .

www.pgadey.ca
@pgadey



UNIVERSITY OF
TORONTO
MISSISSAUGA